



POD-STIK effectively reduces pre-harvest losses caused by 'normal' conditions. Thunderstorms, hailstorms and gales are not 'normal' but 'severe' weather conditions.

To reduce seed loss from pod splitting before and during harvest

An emulsifiable concentrate containing 1 Kg/L Proprietary Surfactant/Polymer Blend

SAFETY INFORMATION

Danger

Causes serious eye damage.

Wear protective gloves/protective clothing/eye protection/face protection.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Immediately call a POISON CENTRE or doctor.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.



Distributed in Ireland by:
Whelehan Crop Protection
Suite 11/12, Bunkilla Plaza, Bracetown Business Park
Clonee, Co. Meath
Telephone: 01 4688900
Fax: 01 8362271



De Sangosse Ltd,
Swaffham Bulbeck,
Cambridge, CB25 0LU
Tel: 01223 811215
Fax: 01223 810020
info@desangosse.co.uk
www.desangosse.co.uk

5L e

DE SANGOSSE



RATE OF USE

POD-STIK is used at a rate of 1.0 L/Ha

MIXING

Three quarter fill the spray tank with water and commence agitation. Add the required quantity of **POD-STIK**. Continue agitation while topping up the tank and during spraying.

DIRECTIONS FOR USE

POD-STIK is a unique polymer blend which has been specifically developed to dramatically reduce the quantity of seed shed from seed pods splitting before and during harvest.

It does not slow down drying, and has no adverse effects on maturity or quality of seed produced.

POD-STIK may be applied at any time after the pods and seeds are full sized, up to before the pods have started to split - optimally the pods are predominantly green.

Apply in 100-250 L/Ha of water as a medium quality spray.

In specific situations where you plan to use high water volumes please consult your agronomist.

SPRAYER CLEANOUT

POD-STIK must not be allowed to dry out in the sprayer, spray lines or nozzles.

NEVER leave unwashed tanks overnight before cleaning.

Immediately after application rinse the sprayer thoroughly with 0.1% Activator 90 in water and spray out to remove any **POD-STIK** residues from the sides of the tank, pump, spray lines and nozzles.

If **POD-STIK** is to be applied the next day, fill tank with water, spray out sufficient to flush pump, spray lines and nozzles, then top up with water.

Provided the **POD-STIK** does not dry out, it will not cause blockages.

For Specific Recommendations contact De Sangosse Ltd or its appointed distributors.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their storage, mixing or use, all conditions and warranties, statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

POD-STIK is a Registered Trademark of De Sangosse Ltd.