



*Adjuvant technology to improve the performance of many herbicides and their tank mix partners*

An emulsifiable concentrate formulation containing 210 g/kg alkoxylated tallow amine, 380 g/kg alcohol ethoxylates, 75 g/kg natural fatty acids and 210 g/kg polyalkylene glycol.

**FOR USE ONLY AS AN AGRICULTURAL/  
HORTICULTURAL/ FORESTRY WETTING AGENT**

**ADJ 0541**

**PCS Number 92243**



**HARMFUL IF SWALLOWED  
IRRITATING TO SKIN  
RISK OF SERIOUS DAMAGE  
TO EYES  
HARMFUL TO AQUATIC  
ORGANISMS, MAY CAUSE  
LONG-TERM ADVERSE EFFECTS  
IN THE AQUATIC ENVIRONMENT**

**HARMFUL**

**Protect From Frost**

**Contents: 1 Litre €**



**DE SANGOSSE**

Distributed in Ireland by Wleehan Crop Protection, Finglas,  
Dublin 11, telephone 01 8068600

De Sangosse Ltd.,  
Swaffham Bulbeck,  
Cambridge, CB25 0LU.  
Tel: 01223 811215  
Fax: 01223 810020  
info@desangosse.co.uk  
www.desangosse.co.uk



**STATUTORY CONDITIONS  
RELATING TO USE**

FOR USE ONLY AS AN AGRICULTURAL / HORTICULTURAL /  
FORESTRY WETTING AGENT

Maximum Concentration: 0.1% Spray solution.

**READ ALL PRECAUTIONS BEFORE USE. ADJ 0541**



**PRECAUTIONS**

The (COSHH) Control of Substances Hazardous to Health  
Regulations may apply to the use of this product at work

**OPERATOR PROTECTION**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:  
**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS),  
SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION  
(FACESHIELD) when handling the concentrate. However engineering  
controls may replace personal protective equipment if a COSHH assessment  
shows that they provide an equal or higher standard of protection.**

**HARMFUL IF SWALLOWED**

**IRRITATING TO SKIN**

**RISK OF SERIOUS DAMAGE TO EYES**

**IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY** with plenty of water and seek medical advice.

**AFTER CONTACT WITH SKIN, WASH IMMEDIATELY** with plenty of water. **WHEN USING, DO NOT EAT, DRINK OR SMOKE.**

**IF YOU FEEL UNWELL,** seek medical advice (show label where possible).

Operators must follow all operator protection instructions for the pesticide. Where the personal protective equipment specified on the adjuvant label is different to that specified on the pesticide label, the highest level of protective equipment must be worn.

**ENVIRONMENTAL PROTECTION**

**USE APPROPRIATE CONTAINER** to avoid environmental contamination  
**DO NOT USE** with any pesticide which is to be applied in or near water.  
**HARMFUL** to aquatic organisms, may cause long-term adverse effects in the aquatic environment

**DO NOT CONTAMINATE SURFACE WATER OR DITCHES** with chemical or used container

**DO NOT EMPTY INTO DRAINS;** dispose of this material and its container in a safe way

**STORAGE AND DISPOSAL**

**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.**  
**KEEP OUT OF REACH OF CHILDREN**

**WASH OUT CONTAINER THOROUGHLY,** and dispose of safely.

**THE CONTAINER MUST NOT BE RE-USED** for any purpose.

**THIS MATERIAL AND ITS CONTAINER** must be disposed of as hazardous waste

**TO AVOID RISKS TO MAN AND THE ENVIRONMENT,  
COMPLY WITH THE INSTRUCTIONS FOR USE.**



**DIRECTIONS FOR USE**

Torpedo-II may be used at a maximum concentration of 0.1% spray solution on the following crops, up to the growth stages specified, with all approved pesticides applied up to their full approved rate.

Crop	Growth Stages before which Torpedo-II may be used with approved pesticides
Cereals	up to and including GS 52 (1/4 inflorescence)
Oilseed rape	up to and including 10% potential pods
Linseed	up to and including 10% capsules formed
Sugar beet	up to and including 6 fully expanded leaves
Potatoes	up to and including tuber initiation

In addition to the above conditions Torpedo-II is authorised for use with all approved pesticides on all edible crops provided that the statutory conditions of use for the pesticide are followed and that the pesticide is not used at more than 50% of the maximum approved rate for that application.

Torpedo-II is also authorised for use with all approved pesticides on all non-edible crops provided that the statutory conditions of use for the pesticide are followed. Torpedo-II may be used to reduce the drift of sprays where this is a concern. Dow AgroSciences LLC recommend Torpedo-II as an approved adjuvant with BROADWAY STAR.

For other Specific recommendations contact De Sangosse Ltd. or its appointed Distributors.



**RATES OF USE**

Torpedo-II is authorised for use up to a maximum concentration of 0.1% dilution of the final water volume e.g:-

Water Volume	Required Quantity of Torpedo-II @ 0.1%
200 litres/ha	200 ml
400 litres/ha	400 ml



**MIXING INSTRUCTIONS**

Add the bulk of the water to the spray tank and commence agitation. Add the recommended quantity of pesticide followed by the required quantity of Torpedo-II. Continue agitation while topping up the tank and during spraying.

**CONDITIONS OF SUPPLY**

All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their storage, mixing or use, all conditions and warranties, statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

*Torpedo-II* ® is a trademark of De Sangosse Ltd. All other Trademarks acknowledged.